

Reply to the Letter: Knowledge of the Prevalence and Epidemiology of Autoimmune Hepatitis

Gabriel Sebastián Díaz-Ramírez,^{1*}  Diego Jiménez,²  Diana Escobar,³  Carlos Vargas,⁴  Carlos Rojas,⁵  Nelson Rojas.⁶ 

OPEN ACCESS

Citation:

Díaz-Ramírez GS, Jiménez D, Escobar D, Vargas C, Rojas C, Rojas N. Reply to the Letter: Knowledge of the Prevalence and Epidemiology of Autoimmune Hepatitis. *Revista. colomb. Gastroenterol.* 2023;38(3):396. <https://doi.org/10.22516/25007440.1128>

¹ Hepatologist, Fundación Valle del Lili. Cali, Colombia.

² Hepatologist, head of the hepatology service, Fundación Valle Del Lili. Cali, Colombia.

³ Hepatologist, Fundación Valle del Lili. Cali, Colombia.

⁴ Internal medicine resident, Universidad ICESI. Cali, Colombia.

⁵ Gastroenterologist, Fundación Valle del Lili. Cali, Colombia.

⁶ General practitioner, research assistant, Clinical Research Center, Fundación Valle del Lili. Cali, Colombia.

*Correspondence: Gabriel Sebastián Díaz-Ramírez.
sebastiandiazr@gmail.com

Received: 16/08/2023

Accepted: 18/08/2023



Dear doctors:

Thank you very much for your observations and especially for reading our article with so much interest. Regarding autoimmune hepatitis worldwide, it is more prevalent in adults, 4 to 42 per 100,000 adults, than in the pediatric population, 2.4 to 9.9 per 100,000 children⁽¹⁾. In our study, despite including patients under 16, we did not find patients under 30 who met the inclusion criteria, as referred to in our conclusion. The non-inclusion of patients under 16 in our search is one of the study's limitations, and it is appropriate to highlight it.

REFERENCE

1. Mack CL, Adams D, Assis DN, Kerkar N, Manns MP, Mayo MJ, et al. Diagnosis and Management of Autoimmune Hepatitis in Adults and Children: 2019 Practice Guidance and Guidelines From the American Association for the Study of Liver Diseases. *Hepatology.* 2020;72(2):671-722. <https://doi.org/10.1002/hep.31065>